What is Claimed is:

1. A compound of the formula (I)

$$Ar^{1} \underset{R}{\overset{O}{\underset{}}} \underset{R^{5}}{\overset{}} R^{6}$$

wherein:

5

10

15

25

 Ar^1 is an aromatic carbocycle substituted with one R^1 , and wherein Ar^1 is independently substituted with two R^2 groups and wherein one R^1 and one R^2 on adjacent ring atoms optionally form a 5- or 6-membered carbocyclic or heterocyclic ring;

 R^1 is halogen, NO₂, NH₂, J-N(R^a)-(CH₂)_m- , N(J)₂-(CH₂)_m- , NH₂C(O)-, J-N(R^a)-C(O)-, J-S(O)_m- N(R^a)-, J-N(R^a)-S(O)_m- or heterocycle-(CH₂)_m- wherein the heterocyclic group is optionally substituted by C₁₋₅ alkyl;

 \mathbf{Q} is a N or $\mathbf{CR}^{\mathbf{p}}$;

Y is
$$> CR^pR^v$$
, $-CR^p = C(R^v)$ -, $-O$ -, $-N(R^x)$ - or $> S(O)_m$;

wherein $\mathbf{R}^{\mathbf{a}}$, $\mathbf{R}^{\mathbf{p}}$, $\mathbf{R}^{\mathbf{v}}$, $\mathbf{R}^{\mathbf{x}}$ and $\mathbf{R}^{\mathbf{y}}$ are each independently hydrogen or $C_{1.5}$ alkyl;

X is
$$-CH_2$$
-, $-N(\mathbb{R}^a)$ -, $-O$ - or $-S$ -;

W is N or CH;

each m is independently 0,1 or 2;

J is chosen from C1-10 alkyl and carbocycle each optionally substituted by R^b ;

 R^2 is chosen from C1-6 alkyl, C3-7 cycloalkyl optionally substituted by C1-5 alkyl, C1-4 acyl, aroyl, C1-4 alkoxy, each being optionally partially or fully halogenated, halogen, C1-6 alkoxycarbonyl, carbocyclesulfonyl and -SO₂-CF₃;

5

10

15

each R³, R⁴ and R⁵ are independently chosen from hydrogen, C1-6 alkyl and halogen;

R⁶ is optionally attached at a position *ortho* or *meta* to the N atom of the indicated ring, and is chosen from

a bond, -O-, -O-(CH₂)₁₋₅-, >C(O), -NH-, -C(O)-NH-, -S-, C₁₋₅ alkyl branched or unbranched, C₂₋₅ alkenyl, C₁₋₃ acyl, C₁₋₃ alkyl(OH), heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl and tetrahydrofuranyl, heteroaryl selected from pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, isoxazolyl, thiazolyl, oxazolyl and isothiazolyl or aryl each alkyl, alkenyl, acyl, heterocycle, heteroaryl and aryl are optionally substituted by one to three hydroxy, oxo, C₁₋₃ alkyl, C₁₋₃ alkoxy, C₁₋₅ alkoxycarbonyl, -NR₇R₈ or NR₇R₈-C(O)-;

wherein each R_6 is further optionally covalently attached to groups chosen from:

20

hydrogen, -NR₇R₈, C_{1-3} alkyl, C_{3-6} cycloalkyl C_{0-2} alkyl, hydroxy, C_{1-3} alkoxy, phenoxy, benzyloxy, aryl C_{0-4} alkyl, heteroaryl C_{0-4} alkyl and heterocycle C_{0-4} alkyl, each above-listed heterocycle, heteroaryl and aryl group is optionally substituted by one to three hydroxy, oxo, C_{1-4} alkyl, C_{1-3} alkoxy, C_{1-5} alkoxycarbonyl, NR₇R₈-C(O)- or C_{1-4} acyl;

25

each R_7 and R_8 are independently hydrogen, phenyl C_{0-3} alkyl optionally subtituted by halogen, C_{1-3} alkyl or di C_{1-5} alkyl amino, or R_7 and R_8 are C_{1-2} acyl, benzoyl or C_{1-5} branched or unbranched alkyl optionally substituted by C_{1-4} alkoxy, hydroxy or mono or di C_{1-3} alkyl amino;

30 and

R^b is chosen from hydrogen, C1-5 alkyl, hydroxyC1-5 alkyl, C2-5 alkenyl, C2-5 alkynyl, carbocycle, heterocycle, heteroaryl, C1-5 alkoxy, C1-5 alkylthio, amino, C1-5 alkylamino, C1-5 dialkylamino, C1-5 acyl, C1-5 alkoxycarbonyl, C1-5 acyloxy, C1-5

acylamino, each of the aforementioned are optionally partially or fully halogenated, or $\mathbf{R}^{\mathbf{b}}$ is chosen from C1-5 alkylsulphonylamino, hydroxy, oxo, halogen, nitro and nitrile;

or the pharmaceutically acceptable salts, acids or isomers thereof.

5

10

2. The compound according to claim 1 and wherein:

Y is -O-, -S-, -NH-, -N(CH_2CH_3)- or -N(CH_3)-;

X is -N(**R**^a)-, or -O-; **Q** is CH;

 ${f J}$ is chosen from C1-10 alkyl, aryl or C3-7 cycloalkyl each optionally substituted by ${f R}^b;$

15

25

30

R₂ is independently chosen from C1-6 alkyl, C3-6 cycloalkyl optionally substituted by C1-3 alkyl, acetyl, aroyl, C1-5 alkoxy, each being optionally partially or fully halogenated, halogen, methoxycarbonyl, phenylsulfonyl and -SO₂-CF₃;

20 each R³, R⁴ and R⁵ are hydrogen;

R^b is chosen from hydrogen, C1-5 alkyl, C2-5 alkenyl, C2-5 alkynyl, C3-8 cycloalkylC0-2 alkyl, aryl, C1-5 alkoxy, C1-5 alkylthio, amino, C1-5 alkylamino, C1-5 dialkylamino, C1-5 acyl, C1-5 alkoxycarbonyl, C1-5 acyloxy, C1-5 acylamino, C1-5 sulphonylamino, hydroxy, halogen, trifluoromethyl, nitro, nitrile or R^b is chosen from; heterocycle chosen from pyrrolidinyl, pyrrolinyl, morpholinyl, thiomorpholinyl, thiomorpholinyl sulfoxide, thiomorpholinyl sulfone, dioxalanyl, piperidinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrofuranyl, 1,3-dioxolanone, 1,3-dioxanone, 1,4-dioxanyl, piperidinonyl, tetrahydropyrimidonyl, pentamethylene sulfide, pentamethylene sulfoxide, pentamethylene sulfone, tetramethylene sulfide, tetramethylene sulfoxide and tetramethylene sulfone and heteroaryl chosen from aziridinyl, thienyl, furanyl, isoxazolyl, oxazolyl, thiazolyl, thiadiazolyl, tetrazolyl, pyrazolyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyranyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl,

benzothiazolyl, benzothienyl, quinolinyl, quinazolinyl, naphthyridinyl, indazolyl, triazolyl, pyrazolo[3,4-b]pyrimidinyl, purinyl, pyrrolo[2,3-b]pyridinyl, pyrazolo[3,4-b]pyridinyl, tubercidinyl, oxazo[4,5-b]pyridinyl and imidazo[4,5-b]pyridinyl.

5

10

3. The compound according to claim 2 and wherein:

 Ar^1 is chosen from phenyl, naphthyl, tetrahydronaphthyl, indanyl and indenyl, each Ar^1 is optionally substituted with one R^1 , and independently substituted with two R^2 groups;

Y is -O-, -S- or -N(CH_3)-;

 ${\bf R}^{\bf 6}$ is present, and is chosen from

a bond, -O-, -O-(CH₂)₁₋₅-, -NH-, -C(O)-NH-, C_{1-5} alkyl branched or unbranched, C_{2-5} alkenyl, C_{1-3} alkyl(OH), heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl and tetrahydrofuranyl, or aryl chosen from phenyl and naphthyl, each alkyl, alkenyl, heterocycle and aryl are optionally substituted by one to three hydroxy, C_{1-3} alkyl, C_{1-3} alkoxy, mono or diC_{1-3} alkyl amino, amino or C_{1-5} alkoxycarbonyl;

20

25

wherein each \mathbf{R}_6 is further optionally covalently attached to groups chosen from: hydrogen, -NR₇R₈, C₁₋₃ alkyl, C₃₋₆ cycloalkylC₀₋₂alkyl, hydroxy, C₁₋₃ alkoxy, phenoxy, benzyloxy, phenylC₀₋₄ alkyl, piperazinylC₀₋₄ alkyl, piperidinyl C₀₋₄alkyl, pyrrolidinylC₀₋₄ alkyl, morpholinylC₀₋₄ alkyl, tetrahydrofuranylC₀₋₄ alkyl, triazolyl C₀₋₄alkyl, imidazolyl C₀₋₄alkyl and pyridinyl C₀₋₄alkyl, each abovelisted heterocycle, heteroaryl and phenyl group is optionally substituted by one to three hydroxy, oxo, C₁₋₄ alkyl, C₁₋₃ alkoxy, C₁₋₅ alkoxycarbonyl, -NR₇R₈, NR₇R₈-C(O)- or C₁₋₄ acyl;

30

each $\mathbf{R_7}$ and $\mathbf{R_8}$ are independently hydrogen, phenyl $\mathbf{C_{0.3}}$ alkyl optionally subtituted by halogen, $\mathbf{C_{1.3}}$ alkyl or di $\mathbf{C_{1.5}}$ alkyl amino, or $\mathbf{R_7}$ and $\mathbf{R_8}$ are $\mathbf{C_{1.2}}$ acyl, benzoyl or $\mathbf{C_{1.5}}$ branched or unbranched alkyl optionally substituted by $\mathbf{C_{1.4}}$ alkoxy, hydroxy or mono or di $\mathbf{C_{1.3}}$ alkyl amino.

4. The compound according to claim 3 and wherein:

X is -O-;

5 Y is $-N(CH_3)$ -;

10

15

20

25

30

J is C1-10 alkyl optionally substituted by $\mathbf{R}^{\mathbf{b}}$;

R₂ is independently chosen from C1-6 alkyl, C3-6 cycloalkyl optionally substituted by C1-3 alkyl and C1-5 alkoxy, each being optionally be partially or fully halogenated;

R⁶ is chosen from

a bond, -O-, -O-(CH₂)₁₋₅-, -NH-, -C(O)-NH-, C_{1-5} alkyl branched or unbranched, C_{2-5} alkenyl, C_{1-3} alkyl(OH), heterocycle selected from morpholinyl, piperazinyl, piperidinyl and pyrrolidinyl or phenyl, each alkyl, alkenyl, heterocycle and phenyl are optionally substituted by one to three hydroxy, C_{1-3} alkyl, C_{1-3} alkoxy, mono or diC_{1-3} alkyl amino, amino or C_{1-5} alkoxycarbonyl;

wherein each R_6 is further optionally covalently attached to groups chosen from:

hydrogen, -NR₇R₈, C₁₋₃ alkyl, C₃₋₆ cycloalkylC₀₋₂alkyl, benzyloxy, phenylC₀₋₄ alkyl, piperazinylC₀₋₄ alkyl, piperidinyl C₀₋₄alkyl, pyrrolidinylC₀₋₄ alkyl, morpholinylC₀₋₄ alkyl, triazolyl C₀₋₄alkyl, imidazolyl C₀₋₄alkyl and pyridinyl C₀₋₄alkyl, each abovelisted heterocycle, heteroaryl and phenyl group is optionally substituted by one to three hydroxy, oxo, C₁₋₄ alkyl, C₁₋₃ alkoxy, C₁₋₅ alkoxycarbonyl, amino, NR₇R₈-C(O)- or C₁₋₄ acyl;

each \mathbf{R}_7 and \mathbf{R}_8 are independently hydrogen, phenyl $\mathbf{C}_{0\text{-}2}$ alkyl optionally subtituted by halogen, $\mathbf{C}_{1\text{-}3}$ alkyl or di $\mathbf{C}_{1\text{-}5}$ alkyl amino, or \mathbf{R}_7 and \mathbf{R}_8 are $\mathbf{C}_{1\text{-}5}$ branched or unbranched alkyl optionally substituted by $\mathbf{C}_{1\text{-}4}$ alkoxy, hydroxy or mono or di $\mathbf{C}_{1\text{-}3}$ alkyl amino;

R^b is chosen from hydrogen, C1-5 alkyl, C3-7 cycloalkylC0-2 alkyl, aryl, C1-5 alkoxy, amino, C1-5 alkylamino, C1-3 dialkylamino, C1-3 acyl, C1-5 alkoxycarbonyl, C1-3

acyloxy, C1-3 acylamino, C1-3 sulphonylamino, hydroxy, halogen, trifluoromethyl, nitro, nitrile;

or $\mathbf{R}^{\mathbf{b}}$ is chosen from pyrrolidinyl, pyrrolinyl, morpholinyl, thiomorpholinyl, thiomorpholinyl sulfoxide, thiomorpholinyl sulfone, piperidinyl, piperazinyl,

- 5 piperidinonyl, tetrahydropyrimidonyl, aziridinyl, isoxazolyl, oxazolyl, thiazolyl, thiadiazolyl, tetrazolyl, pyrazolyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl.
- 5. The compound according to claim 4 and wherein:

Ar¹ is formula (A) or (B)

15 wherein:

when Ar^1 is formula (A) then:

 R^1 is NH_2 , J- $N(R^a)$ - $(CH_2)_m$ -, $NH_2C(O)$ -, J- $N(R^a)$ -C(O)-, J- $S(O)_2$ - $N(R^a)$ -, J- $N(R^a)$ - $S(O)_2$ - or heterocycle- $(CH_2)_{1-2}$ - wherein the heterocycle is chosen from pyrrolidinyl, morpholinyl and piperazinyl each optionally substituted by C1-4 alkyl, and

J is C_{1-5} alkyl optionally substituted by $\mathbf{R}^{\mathbf{b}}$;

or

20

when Ar^1 is formula (B) then:

R¹ is hydrogen or halogen;

R₂ is independently chosen from C1-5 alkyl, C3-6 cycloalkyl optionally substituted by C1-3 alkyl and C1-5 alkoxy, each being optionally partially or fully halogenated;

R⁶ is chosen from

a bond, -O-, -O-(CH₂)₁₋₅-, -NH-, -C(O)-NH-, C_{1-5} alkyl branched or unbranched, C_{2-5} alkenyl, C_{1-3} alkyl(OH), heterocycle selected from morpholinyl, piperazinyl, piperidinyl and pyrrolidinyl or phenyl, each alkyl, alkenyl, heterocycle and phenyl are optionally substituted by one to three hydroxy, C_{1-3} alkyl, C_{1-3} alkoxy, mono or diC_{1-3} alkyl amino, amino or C_{1-5} alkoxycarbonyl;

wherein each R₆ is further optionally covalently attached to groups chosen from:

hydrogen, -NR₇R₈, C_{1-3} alkyl, C_{3-6} cycloalkyl C_{0-2} alkyl, benzyloxy, phenyl C_{0-4} alkyl, piperazinyl, piperazinyl C_{1-2} alkyl, piperidinyl, piperidinyl C_{1-2} alkyl, pyrrolidinyl, pyrrolidinyl C_{1-2} alkyl, morpholinyl, morpholinyl C_{1-2} alkyl, triazolyl, triazolyl C_{1-2} alkyl, imidazolyl, imidazolyl C_{1-2} alkyl, pyridinyl and pyridinyl C_{1-2} alkyl, each above-listed heterocycle, heteroaryl and phenyl group is optionally substituted by one to three hydroxy, oxo, C_{1-4} alkyl, C_{1-3} alkoxy, C_{1-5} alkoxycarbonyl, amino, NR_7R_8 -C(O)- or C_{1-4} acyl.

15

10

5

6. The compound according to claim 5 and wherein:

Ar¹ is formula (A) or (B)

$$\mathbb{R}^1$$
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^1
 \mathbb{R}^2
 \mathbb{R}^1

20

and R2 is chosen from

```
when \mathbf{Ar^1} is formula (A) then:
when \mathbf{R^1} is \mathbf{J}\text{-S}(O)_2- \mathbf{N(R^a)}- or \mathbf{J}\text{-N(R^a)}-\mathbf{S}(O)_2- then \mathbf{J} is \mathbf{C_{1-3}} alkyl;
and
when \mathbf{R^1} is \mathbf{NH_2}, \mathbf{J}\text{-N(R^a)}-(\mathbf{CH_2})_{m^-}, \mathbf{NH_2C}(O)-, \mathbf{J}\text{-N(R^a)}-\mathbf{C}(O)-,
or heterocycle-(\mathbf{CH_2})_{1-2}- wherein the heterocycle is chosen from pyrrolidinyl,
morpholinyl, piperazinyl or \mathbf{C1}-4alkylpiperazinyl, then
\mathbf{J} is \mathbf{C1}-3 alkyl optionally substituted by \mathbf{R^b}.
```

7. The compound according to claim 6 and wherein:

R^b is chosen from hydrogen, C1-5 alkyl, C3-6 cycloalkylC0-2 alkyl, phenyl, C1-5 alkoxy, amino, C1-5 alkylamino, C1-3 dialkylamino, C1-3 acyl, C1-5 alkoxycarbonyl, C1-3 acyloxy, C1-3 acylamino, hydroxy, halogen;

- or **R**^b is chosen from morpholinyl, thiomorpholinyl, thiomorpholinyl sulfoxide, thiomorpholinyl sulfone, piperidinyl, piperidinonyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl.
 - 8. The compound according to claim 7 and wherein:

 R^b is chosen from amino, C1-5 alkylamino, C1-3 dialkylamino; or R^b is chosen morpholinyl, piperidinyl and pyridinyl.

- 9. The compound according to claim 6 and wherein:
- 25 Ar^1 is formula (A).

20

- 10. The compound according to claim 6 and wherein: Ar^1 is formula (B).
- 11. The compound according to claim 6 and wherein:Ar¹ is

12. A compound chosen from:

1-Methyl-7-(pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

7-(2-Methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-phenyl)-amide

7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

7-(2-Cyclopropylamino-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

7-[2-(4-Methoxy-benzylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-phenyl)-amide

1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-phenyl)-amide

1-Methyl-7-[2-(2-morpholin-4-yl-ethylamino)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

- 7-[2-(3-Dimethylamino-propylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(3-Dimethylamino-2,2-dimethyl-propylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Dimethylamino-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(6-methyl-2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-pyrrolidin-1-yl-ethylamino)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(piperidin-4-ylamino)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 4-{4-[2-(5-tert-Butyl-3-methanesulfonylamino-2-methoxy-phenylcarbamoyl)-1-methyl-1H-indol-7-yloxy]-pyrimidin-2-ylamino}-piperidine-1-carboxylic acid tert-butyl ester
- 7-{2-[(2-Dimethylamino-ethyl)-methyl-amino]-pyrimidin-4-yloxy}-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxyphenyl)-amide
- 1-Methyl-7-[6-methyl-2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethoxy)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethoxy)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-pyrrolidin-1-yl-ethoxy)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-morpholin-4-yl-ethoxy)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(1-methyl-piperidin-4-yloxy)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethoxy)-6-methyl-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

- 1-Methyl-7-(2-methylcarbamoyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid [5-tert-butyl-3-(2-dimethylamino-ethylcarbamoyl)-2-methoxy-phenyl]-amide
- 7-[2-(2-Dimethylamino-ethylcarbamoyl)-pyridin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid [5-tert-butyl-3-(2-dimethylamino-ethylcarbamoyl)-2-methoxy-phenyl]-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid [5-tert-butyl-2-methoxy-3-(2-morpholin-4-yl-ethylcarbamoyl)-phenyl]-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-carbamoyl-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-methylcarbamoyl-phenyl)-amide
- 1-Methyl-7-(2-vinyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(1,2-Dihydroxy-ethyl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(morpholin-4-ylamino)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-morpholin-4-ylmethyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-morpholin-4-ylmethyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-ylmethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Dimethylaminomethyl-pyridin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-methylcarbamoyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Benzyloxymethyl-pyridin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-morpholin-4-ylmethyl-phenyl)-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid [5-tert-butyl-2-methoxy-3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-dimethylaminomethyl-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (3-amino-5-tert-butyl-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-dibenzylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-methylsulfamoyl-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-[1,3]dioxolan-2-yl-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-methylaminomethyl-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-pyrrolidin-1-ylmethyl-phenyl)-amide
- 1-Methyl-7-{2-[methyl-(1-methyl-piperidin-4-yl)-amino]-pyrimidin-4-yloxy}-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Hydroxymethyl-pyridin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide and
- 1-Methyl-7-(2-methylamino-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid [5-tert-butyl-2-methoxy-3-(2-morpholin-4-yl-ethylamino)-phenyl]-amide
- 1-Methyl-7-(pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-piperazin-1-yl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid [3-methanesulfonylamino-2-methoxy-5-(1-methyl-cyclopropyl)-phenyl]-amide

- 1-Methyl-7-[2-(5-methyl-2,5-diaza-bicyclo[2.2.1]hept-2-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2,5-Diaza-bicyclo[2.2.1]hept-2-yl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert- butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Methoxy-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(4-tert-Butyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-morpholin-4-yl-ethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-{2-[2-(4-methyl-piperazin-1-yl)-ethyl]-pyrimidin-4-yloxy}-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxyphenyl)-amide
- 1-Methyl-7-[2-(2-pyrrolidin-1-yl-ethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethyl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-morpholin-4-yl-ethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-{2-[2-(4-tert-Butyl-piperazin-1-yl)-ethyl]-pyrimidin-4-yloxy}-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(4-tert-Butyl-piperazin-1-ylmethyl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-pyrrolidin-1-ylmethyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2,6-Dimethyl-pyridin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Ethyl-pyridin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

- 1-Methyl-7-[2-(1,2,3,6-tetrahydro-pyridin-4-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Amino-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-pyrrolidin-1-ylmethyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-piperidin-1-ylmethyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-ylmethyl)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-{[(2-dimethylamino-ethyl)-methyl-amino]-methyl}-2-methoxy-phenyl)-amide
- 7-(2-{[(2-Dimethylamino-ethyl)-methyl-amino]-methyl}-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-carbamoyl-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-((1S,4S)-5-methyl-2,5-diaza-bicyclo[2.2.1]hept-2-yl)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-[1,4]diazepan-1-yl)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-[1,4]Diazepan-1-yl-pyridin-4-yloxy)-1-methyl-1-H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-piperazin-1-yl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-trideuterio-7-(2-piperazin-1-yl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(Hexahydro-pyrrolo[1,2-a]pyrazin-2-yl)-pyridin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-trideuterio-7-[2-(4-methyl-piperazin-1-yl)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid [3-methanesulfonylamino-2-methoxy-5-(1-methyl-cyclopropyl)-phenyl]-amide

- 7-[2-((S)-3-Dimethylamino-pyrrolidin-1-yl)-pyridin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid [3-methanesulfonylamino-2-methoxy-5-(1-methyl-cyclopropyl)-phenyl]-amide
- 7-[2-((S)-3-Dimethylamino-pyrrolidin-1-yl)-pyridin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid [3-methanesulfonylamino-2-methoxy-5-(1-methyl-cyclopropyl)-phenyl]-amide
- 1-Methyl-7-[2-(4-methyl-piperazine-1-carbonyl)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide and
- 1-Methyl-7-[2-(piperazine-1-carbonyl)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

or the pharmaceutically acceptable salts, acids or isomers thereof.

13. A compound chosen from:

- 7-(Pyrimidin-4-yloxy)-benzo[b]thiophene-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(Pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(Pyrimidin-4-yloxy)-benzofuran-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(pyrimidin-4-ylsulfanyl)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(pyrimidin-4-ylamino)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(pyridin-3-yloxy)-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Benzylamino-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-{2-[(pyridin-2-ylmethyl)-amino]-pyrimidin-4-yloxy}-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2-Imidazol-1-yl-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-[1,2,3]triazol-1-yl-ethylamino)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

- 7-[2-(3-Dimethylamino-propylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid [2-methoxy-5-(2,2,2-trifluoro-1-trifluoromethyl-ethyl)-phenyl]-amide
- 7-{2-[(2-Dimethylamino-ethyl)-methyl-amino]-pyrimidin-4-yloxy}-1-methyl-1H-indole-2-carboxylic acid (4-chloro-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-[2-(4-Acetyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (4-chloro-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-[2-(2-Dimethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (2-methoxy-5-trifluoromethoxy-phenyl)-amide
- 7-[2-(4-Dimethylamino-piperidin-1-yl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(3-Dimethylamino-pyrrolidin-1-yl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(1-methyl-piperidin-4-ylamino)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(1-Acetyl-piperidin-4-ylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(2-morpholin-4-yl-ethoxy)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-[2-(2-Imidazol-1-yl-ethoxy)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(2-Imidazol-1-yl-ethoxy)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (4-chloro-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-methylcarbamoyl-phenyl)-amide
- 7-(2-Amino-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-carbamoyl-2-methoxy-phenyl)-amide
- 7-(2-Amino-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid [5-tert-butyl-3-(2-dimethylamino-ethylcarbamoyl)-2-methoxy-phenyl]-amide

- 7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-dimethylaminomethyl-2-methoxy-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-pyrrolidin-1-ylmethyl-phenyl)-amide
- 7-[2-(2-Dimethylamino-ethylamino)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-2-methoxy-3-morpholin-4-ylmethyl-phenyl)-amide
- 1-Methyl-7-(2-morpholin-4-ylmethyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (4-chloro-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-[2-(3-Dimethylamino-pyrrolidin-1-ylmethyl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-(2-Carbamoyl-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-(2-morpholin-4-ylmethyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (2-methoxy-3-morpholin-4-ylmethyl-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (2-methoxy-3-morpholin-4-ylmethyl-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-(2-morpholin-4-ylmethyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-(1'-tert-Butyl-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-(2-methylaminomethyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-(2-pyrrolidin-1-ylmethyl-pyridin-4-yloxy)-1H-indole-2-carboxylic acid (2-methoxy-3-morpholin-4-ylmethyl-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-[2-(2-morpholin-4-yl-ethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (3-dimethylaminomethyl-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-(2-pyrrolidin-1-ylmethyl-pyrimidin-4-yloxy)-1H-indole-2-carboxylic acid (2-methoxy-3-pyrrolidin-1-ylmethyl-5-trifluoromethyl-phenyl)-amide
- 7-(2-Dimethylaminomethyl-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid [2-methoxy-3-(4-methyl-piperazin-1-ylmethyl)-5-trifluoromethyl-phenyl]-amide

- 7-(2-Dimethylaminomethyl-pyridin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-(2-Dimethylaminomethyl-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-ylmethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 7-(2-Dimethylaminomethyl-pyrimidin-4-yloxy)-1-methyl-1H-indole-2-carboxylic acid (3-methanesulfonylamino-2-methoxy-5-trifluoromethyl-phenyl)-amide
- 1-Methyl-7-[2-(2-morpholin-4-yl-ethyl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid [3-methanesulfonylamino-2-methoxy-5-(1-methyl-cyclopropyl)-phenyl]-amide
- 1-Methyl-7-[2-(1-methyl-piperidin-4-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 7-[2-(1-Cyclopropyl-piperidin-4-yl)-pyrimidin-4-yloxy]-1-methyl-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide
- 1-Methyl-7-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yloxy]-1H-indole-2-carboxylic acid (3-dimethylaminomethyl-2-methoxy-5-trifluoromethyl-phenyl)-amide and
- 1-Methyl-7-[2-(1-methyl-pyrrolidin-3-ylamino)-pyridin-4-yloxy]-1H-indole-2-carboxylic acid (5-tert-butyl-3-methanesulfonylamino-2-methoxy-phenyl)-amide

or the pharmaceutically acceptable salts, acids or isomers thereof.

- 14. A pharmaceutical composition containing a pharmaceutically effective amount of a compound according to claim 1 and one or more pharmaceutically acceptable carriers and/or adjuvants.
- 15. A method of treating an oncological disease comprising administering to a patient a pharmaceutically effective amount of a compound according to claim 1.
 - 16. A method of treating a disease or condition chosen from osteoarthritis, atherosclerosis, contact dermatitis, bone resorption diseases, reperfusion injury, asthma, multiple sclerosis, Guillain-Barre syndrome, Crohn's disease, ulcerative colitis,

5

10

20

25

psoriasis, graft versus host disease, systemic lupus erythematosus, insulin-dependent diabetes mellitus, rheumatoid arthritis, toxic shock syndrome, Alzheimer's disease, diabetes, inflammatory bowel diseases, acute and chronic pain, stroke, myocardial infarction alone or following thrombolytic therapy, thermal injury, adult respiratory distress syndrome (ARDS), multiple organ injury secondary to trauma, acute glomerulonephritis, dermatoses with acute inflammatory components, acute purulent meningitis, syndromes associated with hemodialysis, leukopherisis, granulocyte transfusion associated syndromes, necrotizing entrerocolitis, restenosis following percutaneous transluminal coronary angioplasty, traumatic arthritis, sepsis, chronic obstructive pulmonary disease and congestive heart failure, said method comprising administering to a patient a pharmaceutically effective amount of a compound according to claim 1.

- 17. A method of treating a disease or condition requiring anticoagulant or fibrinolytic therapy, said method comprising administering to a patient a pharmaceutically effective amount of a compound according to claim 1.
 - 18. A process of making a compound of the formula (I):

$$Ar^{1} \underset{H}{\overset{O}{\bigvee}} \underset{R^{3}}{\overset{V}{\bigvee}} \underset{R^{4}}{\overset{V}{\bigvee}} \underset{R^{4}}{\overset{R^{6}}{\bigvee}}$$

Ar₁, X, Y, Q, W, R³, R⁴, R⁵, R⁶ and R^y are defined in claim 1; said process comprising

coupling under suitable conditions an amine bearing Ar¹ carboxylic acid of the formula (III), where P is a protecting group,

removing the protecting group P to provide an intermediate of formula (V) under suitable conditions;

coupling under suitable conditions the intermediate (V) with a halo heterocycle VI (Z = halogen) bearing R^6 in the presence of a suitable base to provide a compound of the formula (I):